

Hop Botany, Cultivation, and Breeding

Jason Perrault
Perrault Farms, Inc.
Hop Breeding Company, LLC
Select Botanicals Group, LLC-John I. Haas, Inc.

Perrault Farms, Inc.
Yeggenia, WA



SELECT
BOTANICALS
G-R-O-U-P

Hop Botany, Cultivation, and Breeding

- Importance of hops.
- Basic botanical information.
- Crop development and cultivation.
- Impact of hop varieties.
- Variety development.



The Importance of Hops



Regional Economic Importance



- U.S. Production centered in the PNW.
 - 77% in WA.
 - 16% in OR.
 - 7% in ID.
- 2011 value (US) = \$180 million
- Annually Top 12 in crop value for Washington

Humulus spp. Overview

- Family:
Cannabaceae
 - Cannabis*
 - C. sativa*
 - Humulus*
 - H. japonicus*
 - H. yunnanensis*
 - H. lupulus*



(Neve 1991)

Humulus lupulus

- "Hops"
- Dioecious, perennial, climbing vine
- Indigenous to the Northern Hemisphere
 - Origins in Europe:
 - H. lupulus* var. *lupulus*
 - Origins in Asia (mainly Japan):
 - H. lupulus* var. *cordifolius*
 - Origins in North America:
 - H. lupulus* var. *pubescens*
 - H. lupulus* var. *neomexicanus*
 - H. lupulus* var. *lupuloides*

Hop Basics


- Dioecious (male and female plants).
 - Genetically complex.
 - Male-no commercial value
 - Female-Produces the valued strobiles, "cones"
- Annual above ground.
- Perennial below.
 - Allows for clonal propagation.
- Climbing bine requiring a support system.
- Photoperiod sensitive

Dioecious Plants

- Separate male and female plants
- Commercial value derived from the strobiles or "cones" of the female plant
- Male plants utilized only for hybridization
- Pollination results in:
 - Unwanted seeds
 - Increased cone size

The "Cones"

- These are the manufacturing unit of the commercial hop plant.
 - The cones contain lupulin glands (actually modified vine hairs).
 - These glands contain the chemistry we are after:
 - Essential oils: well over 100 compounds, contribution to aroma.
 - Soft resins: beta acids, and the all important alpha acids.
 - Lupulin accounts for 20 – 30 % of cone weight.



Mature Female "Cones"

Male flowers at anthesis


Annual vs. Perennial Growth

- The above ground portion of the stem is annual.
 - Dies off at dormancy.
- The root is perennial, can survive low winter temps.
 - Requires a dormant period.
- The plant also produces rhizomes (below ground stems).
 - Buds become new spring growth.
 - Easily propagated from cuttings.



Clonal Propagation

- Propagation of hops purely vegetative
 - Root cuttings
 - Layering
 - Softwood cuttings
- Resulting plants genetically identical to parent material



Climbing Bines

- In the wild-usually found climbing on companion species. In cultivation, trellis is used.
- Typical Field Setup:
 - Trellis 18' high
 - Plant spacing at 3.5' x 14' or 7' x 7'.
 - Result is 889 plants per acre
 - Anchored twine is used to support plant growth.
- The vine wraps clockwise around string.
 - Function of phototropism and thigmotropism (Light and Touch).
- Rapid growth: The hop plant will grow a foot or more a day under ideal conditions. 18-25' in a season.

Photoperiod Sensitive

- Hops are a short day plant.
 - Under a critical number of light hours - floral initiation.
 - Also node dependant.
 - Over the critical amount, vegetative growth.
 - In shorter day areas, flowering occurs as soon as the node requirement in met-yield not maximized.
 - In longer day areas-vegetative growth is maximized prior to shortening days of mid to late summer.
- Results in defined "Production Stages"

Developmental Physiology of the Hop Plant (or Production Stages)

- The hop plant goes through numerous stages of growth throughout the year.
 - Each stage has its own unique characteristics.
 - Therefore each stage of growth requires its own unique management scheme.
- Main Stages of Growth
 - Dormancy
 - Spring regrowth
 - Vegetative Growth
 - Reproductive Growth
 - Preparation for Dormancy

Dormancy: October through February

- October through February:
 - Late summer the plant allocates photosynthetically derived starches to storage roots
 - The starch is converted into soluble sugars.
 - These sugars are the energy needed to commence spring regrowth.

Dormancy: October through February

- What's going on in the field? Not a whole lot.
 - Compost applications.
 - Working the ground.
 - Prepping new yards.

Spring Regrowth March through May

- The end of dormancy is signaled by increasing day length and increasing temperatures in the spring.
 - The plant utilizes the soluble sugars as energy to emerge from dormancy and commence regrowth.
 - The initial regrowth occurs rapidly producing vines unsuitable for crop production.
 - The plant relies on the energy reserves of the root until the end of May, at which time the starches and soluble sugars reach their lowest points of the year.
 - To maximize plant health, supplemental nutrient management will be needed.

Spring Regrowth March through May

- What's happening in the field?
 - Spring pruning- March-April
 - Effort to maximize consistency for training
 - Weed control
 - Applications of dry fertilizer
 - Twining
 - Training- one of the most important aspects of crop production.
 - Timing is varietal specific and critical.
 - Generally target 3 vines per string.
 - Irrigation begins

Importance of Photoperiod Sensitivity

- Hops are a short day plant.
 - Under a critical number of light hours (more accurately it is the length of the dark period)-floral initiation.
 - Also node dependant.
 - Over the critical amount, vegetative growth.
 - In shorter day areas, flowering occurs as soon as the node requirement in met-yield not maximized.
 - In longer day areas-vegetative growth is maximized prior to shortening days of mid to late summer.

Vegetative Growth


- The vegetative growth stage, for the purposes of crop production, occurs from the end of May through the end of July.
- It can be separated into two phases:
 1. From May to the end of June/early July: Plant growth is mainly found in the main vine and leaves.
 2. July: The bulk of the above ground growth occurs in lateral production.

Vegetative Growth

- This is a critical period:
 - The plants reserves are used up.
 - The plant, even now, is already determining how much it is going to yield.
 - We need to manage plant health aggressively during this stage of growth.
 - The goal should be to maximize the health of the plant, while managing growth-this is tricky.

Vegetative Growth

- What's Happening in the field?
 - Monitor, monitor, monitor
 - Pest/Disease/Weed control
 - Irrigation
 - Fertility



Preparation for Dormancy:

End of August to beginning of September:

- While not really a stage of growth, it is important in the development of the crop for next year.
 - Photosynthetic production of carbohydrates exceeds the needs of plant development.
 - The excess is transported to the roots for storage in the form of starch.
 - Both the dry weight of the roots as well as starch content has peaked by October.
 - The shortening days of late summer signal this transition, followed by cold October temperatures- Dormancy starts.






Preparation for Dormancy:

End of August to beginning of September:

- What's Happening in the field?
- Harvest commences.

Harvest

- Vines are cut and transported to picker.
 - Alternatively, use field strippers
- Material is ran through stationary machine, cones are separated.
- Cones dried for 8-12 hours to 10% moisture.
- Dried cones are cooled (ambient) for 12 to 24 hours.
- Baled and transported immediately to cold storage.



How important is this?

- Hop Supply Chain: Each link on the supply chain affects subsequent links.
 - The efficiency of a hop has a corresponding impact on the chain.



Developing Objectives

- The hop trade consists of two distinct markets:
 - Alpha/Bitter
 - Processed hops.
 - Yield measured in Kg. Alpha per acre.
 - Typically high alpha varieties, increasingly aroma.
 - Aroma
 - Minimal processing.
 - Yield measured in lb. acre.
 - Typically aroma varieties, some high alphas.
- This is an important consideration when setting objectives.

Specific Objectives

- High yielding high alpha cultivars.
 - Super Varietal
- High yielding aroma cultivars.
 - Improvements on the classics
 - Specialty / dual purpose
 - Organic
- Goal is to combine the above with:
 - Pest and disease resistance.
 - Good storage stability.
 - Desirable brewing characteristics (i.e. low cohumulone, specific oil components).

Parental Selection

- Remember- Hops are dioecious.
 - Distinct male and female plants.
 - Obligate out-crossers, cannot self pollinate.
 - High level of diversity (heterozygosity).
 - Hybrid vigor (Heterosis).
 - Seed propagation not possible.
 - Easily clonally propagated- traits can be "fixed" in single generation.
 - Each new variety results from a single plant.
 - Millions from one.

Crossing



Left: Collection of male flowers for isolation of pollen.
Above: Application of pollen to a bagged receptive female.

From Crosses to seedlings



Typically start with 20,000 - 50,000 genotypes in any given year.




Seedlings are screened in the greenhouse for Powdery Mildew, then planted to the field.



Typically eliminate 75 - 90% of the starting population.


Hop Breeding Scheme



Cultivar Release: Year 11

- After 8 - 10 years of evaluation, release is considered.
 - Private varieties: PVP begins.
- The work is far from over, success is dependant on:
 - Continued agronomic success.
 - Grower acceptance, usually short term.
 - Brewer acceptance, long term.

Yields of New U. S. Aroma Varieties



Organic Hop Breeding: YCR 4, Palisade® Example


Conventional




Organic



Organic Hop Breeding: YCR 4, Palisade® Example



Organic Yield Comparison: Top Ten Breeding Lines vs. Palisade




Future Trends in Hop Breeding

- Molecular research
 - Marker assisted selection
 - Gene mapping
 - Gene functionality
- Non-brewery usage
- Continuing conversion to new varieties
 - Driven by disease pressure, storage issues, basic economic pressures, and continued growth in craft brewing.
 - Increases focus on AROMA

Parting Thoughts: Overcoming Challenges

- Do your homework.
 - Know your plant, environment.
 - Know your market.
 - Organic? Local? Sustainability?
 - Hops as a commodity, does not work.
- Developing relationships is key.



Conclusion

- Hops are complex, high cost crop.
 - Not necessarily high value.
 - Knowledge of the growth stages is critical.
- Hop breeding is a necessary, functional step in the hop supply chain.
 - Supplies the varieties which decrease costs in subsequent steps.
 - It is a long complex process which demands commitment.
- Marketing is critical.

