

# FIFRA Section 24(c) Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Delegate<sup>®</sup> WG

EPA Reg. No. 62719-541

EPA 24(c) Special Local Need SLN No. MI-170002

### One-day Pre-harvest Interval in Bushberries (Subgroup 13-07B<sup>1</sup>) Except Lingonberry

For Distribution and Use Only Within the State of Michigan

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2022.

#### Directions for Use

##### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Delegate<sup>®</sup> herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Delegate is subject to all use precautions and limitations imposed by the label affixed to the container for Delegate.

Refer to product label for Delegate for General Use Precautions, Mixing Directions, and Application Directions.

**Application Timing:** Treat using local extension recommendations and action thresholds. Heavy infestations may require repeat applications, but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** The amount of Delegate WG applied per acre will depend upon plant size, volume of foliage present and pest pressure. Use a lower rate in the rate range for light infestations and/or small plants and a higher rate in the rate range for heavy infestations and/or larger plants.

#### Pests and Application Rates:

Pests	Delegate WG (oz/acre)
armyworms	3 – 6
blueberry gall midge (suppression)	
blueberry maggot (suppression)	
cherry fruitworm	
cranberry fruitworm	
currant fruit fly (suppression)	
European grapevine moth	
fireworms	
gypsy moth	
leafrollers	
light brown apple moth	
loopers	
thrips (suppression)	

**Resistance Management:** Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Do not make more than three applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative,

extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

**Special Use Restrictions for Preharvest Interval of one day:**

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- Do not apply more than a total of 17.9 oz of Delegate WG (0.281 lb ai spinetoram) per acre per year.
- **Maximum Number of Applications:** Do not make more than three applications per calendar year. See Resistance Management regarding number of applications for specific pests. See Resistance Management and Special Use Restrictions regarding number of applications for specific pests and locations.
- **Minimum Treatment Interval:** Do not make application less than 6 days apart between the first and second applications. Do not make applications less than 12 days apart between the second applications and the third application. The third application is applied at 1 day of harvest.

®Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

R359-034

Accepted: 06/07/17

Replaces: Initial printing