

Fungicide(s)										
Active ingredient	Product	Rate/A (fl. oz)	Powdery mildew	leaf/ glume blotch	Septoria leaf blotch	Stripe rust	Leaf rust	Stem rust ⁴	Head scab	Harvest Restriction
Metconazole 8.6%	Caramba 0.75 SL	10.0 - 17.0	VG ¹	VG	-- ²	E	E	E	G	30 days
Propiconazole 41.8%	Tilt 3.6 EC PropiMax 3.6 EC Bumper 41.8 EC	4.0	VG	VG	VG	VG	VG	VG	P	Feekes 10.5
Prothioconazole 41%	Proline 480 SC	5.0 - 5.7	-- ²	VG	VG	-- ²	VG	VG	G	30 days
Tebuconazole 38.7%	Folicur 3.6 F ³	4.0	G	VG	VG	E	E	E	F	30 days
Prothioconazole 19% Tebuconazole 19%	Prosaro 421 SC	6.5 - 8.5	G	VG	VG	E	E	E	G	30 days

¹ Efficacy categories: NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent.

² Insufficient data to make statement about efficacy of this product.

³ Multiple generic products containing tebuconazole may also be labeled in some states. These products include: Muscle 3.6 F, Orius 3.6 F, Tebucon 3.6 F, Tebustar 3.6 F, Tebuzol 3.6 F, Tegrol, & Toledo.

⁴ Estimates of fungicide efficacy against stem rust are based on a small number of observations and may be less reliable than the ratings for other diseases.

Source: Management of Fusarium Head Blight (Scab), Diane Brown-Rytlewski, Martin Nagelkirk, and William Kirk, MSU Extension.

Reviewed by William Kirk and Janet Lewis, MSU.